

A stability analysis of radiative shocks in the presence of a transverse magnetic field

Babulakshmanan Ramachandran^{*} & Michael D. Smith[†]

Armagh Observatory, College Hill, Armagh BT61 9DG, Northern Ireland

Accepted Received ; in original form

ABSTRACT

Radiative shock waves may be subject to a global thermal instability in which the cooling layer and shock front undergo growing resonant oscillations. For strong hydrodynamic shocks, the presence of the overstability depends on the temperature and density indices of power-law cooling functions and the specific heat ratio, α , β and γ , respectively. Here, we investigate the stabilising influence of a transverse magnetic field by introducing the shock Alfvén number, M_a as a fourth parameter. We thus investigate the stability criteria for both molecular and atomic shocks under a wide range of conditions. In particular, we find that all molecular shocks in which the cooling increases with the temperature ($\alpha > 0$) are stabilised to the first four modes if $M_a < 20$ ($\beta = 2$). For $\alpha = -0.5$, the first overtone remains stable only for $M_a < 8$. We conclude that molecular shocks in the interstellar medium are probably stabilised by a transverse magnetic field unless exceptional circumstances arise in which the cooling strongly increases as the gas cools.

Key words: shock waves – MHD – magnetic fields – instabilities – ISM: molecules

1 INTRODUCTION

To interpret observations of an astrophysical shock we need to know if it can be compared to steady state models or if it is subject to an instability. In fact, the cooling immediately downstream of a shock front may lead to an overstability of the entire radiative shock wave (?). For hydrodynamic flows, a linear analysis yields conditions for growing oscillations as well as their particular frequencies (?), and many analytical and numerical studies have been presented, as recently summarised by (?; hereafter, Paper 1). However, the growth of the oscillations will be damped by a magnetic field. A transverse field not only reduces the immediate post shock temperature and compression but could also completely stabilise the cooling layer. The first question is: how strong does the field have to be to ensure stability? This question has been answered for the important case of fast shocks into atomic gas (??). Here, we extend these results to include power-law cooling functions pertaining to other interstellar conditions: a molecular gas and a medium in which the energy levels relevant to the cooling are maintained in local thermodynamic equilibrium. The conclusions take on added significance through the latest campaigns to explore infrared and submillimetre regimes with spectroscopic methods (e.g. with Spitzer and Herschel).

A linear analysis of plane-parallel radiative shocks with a transverse field was performed by ?. They assumed a specific heat ratio (γ) of 5/3 and a cooling function (Λ) $\propto \rho^2 T^\alpha$, where ρ is the density and T is the temperature. Only the index α determines the stability in the absence of a magnetic field provided the shock is strong. The latter assumption eliminates the Mach number, M , from the problem. Here, $M = u_{in}/c_s$ where u_{in} is the shock speed and c_s is the upstream sound speed.

On the other hand, the Alfvén number $M_a = u_{in}/v_a$, where v_a is the upstream Alfvén speed, is a second variable when a significant magnetic field is present. (?; hereafter TD93) found, as an example, that even a quite weak field ($M_a < 8$) will suppress the growth of all modes of oscillation for $\alpha > 0$. For $\alpha = 0.5$, even weaker fields ($M_a < 33$) are enough to stabilise a shock. Their numerical study agreed with the linear analysis and also revealed that for shock speeds $v_s < 160 \text{ km s}^{-1}$, radiative shocks occurring in interstellar gas with $n_H \leq 0.4 \text{ cm}^{-3}$ may be magnetically stabilised. Further simulations by ? however, also demonstrated that the typical interstellar field may not be sufficient to stabilise shocks if $\alpha < 0$. The fundamental mode can generate very large amplitude oscillations. Moreover, even when the fundamental is stabilised, the overtones can still produce higher frequency oscillations of substantial amplitude.

The general linear analysis for hydrodynamic flows was extended in Paper 1 to include molecular shocks. The dependence on three parameters was considered: α , β and γ

^{*} E-mail: brc@arm.ac.uk

[†] E-mail: mds@arm.ac.uk

where β is the density dependence of the cooling:

$$\Lambda \propto \rho^\beta T^\alpha. \quad (1)$$

A strong dependence on γ was found, with the regime of overstability significantly reduced for molecular shocks. In particular the fundamental mode grows only for $\alpha < -0.24$ in the molecular equivalent ($\gamma = 7/5$ and $\beta = 2$) of the atomic case (for which $\alpha < 0.38$ is required). However, the overtones are more significant in molecular shocks with, for example, the first overtone growing for $\alpha < 0.66$. Building on this work, we introduce the Alfvén number, here as the fourth parameter, to determine the influence of the magnetic field.

In the molecular shocks discussed here, we presume that the fraction of mass in ions is sufficiently high so that ion-magnetosonic waves cannot propagate ahead of the shock front. Otherwise, if the fraction of ions would be low, the magnetic field will drift through the neutral molecules, a process termed ambipolar diffusion. The friction between the ions and neutrals may then replace the molecular viscosity, changing the nature of the shock front. We discuss this restriction in § 3. Other dynamic instabilities are then possible in such Continuous Shocks (as well as in Jump Shocks, see Paper 1 for a summary). The Jump Shocks studied here thus require a source of ionisation that does not cause wholesale molecule dissociation. This may occur near the edges of molecular clouds partially exposed to the ultraviolet from nearby OB associations, as proposed for the HH 90/91 shock (?). The ions can be dust grains, molecules or atoms, according to the physical conditions.

In this paper, we extend the formalism of TD93. Starting with the steady state radiative shock structure, we perturb the shock velocity and determine the growth or damping rate in addition to the oscillatory period. We study the one dimensional case while ignoring thermal conduction as well as radiation transfer. In the hydrodynamic case, a stationary wall of infinite density is assumed to lie downstream of the shock front. In the MHD analysis, the density in the downstream gas rises towards a finite constant value. In order to make the problem tractable, we not only ensure that the temperature continues to fall, reaching zero Kelvin, but that the temperature perturbation also vanishes at the downstream boundary. Then, waves escaping downstream are constrained to be pure Alfvén waves, leading to a straightforward boundary condition. In this respect, it should be noted that the published versions of equation (A27) of TD93 (where the minus sign should be replaced by a +) and equation (2) of ? (where the conditionals should be reversed) are incorrect.

In Paper 1, we undertook a linear stability analysis similar to ? while taking a general γ so as to encompass molecular shocks. By also generalising the density index, we take into account different physical situations. In particular, $\beta = 1$ corresponds to cases where the energy levels of the atoms or molecules which provide the dominant cooling are in local thermodynamic equilibrium. For example, this can correspond to H_2 cooling in warm gas for densities above $\sim 10^4 \text{ cm}^{-3}$ provided hydrogen atoms act as the main collision partner. The specific heat ratio $\gamma = 5/3$ remains appropriate for dissociative shocks. The value $\gamma = 7/5$ applies to a pure H_2 gas while, more realistically for the interstellar




Figure 1. A sketch of the steady shock configuration. The shock jump conditions are defined by equations 16, 17 and 18; the total compression, $-w_f$, is given by equation 22. Note that under strong shock conditions, the upstream thermal pressure is equated to zero.

medium, $\gamma = 10/7$ accounts for the addition of ten per cent helium atoms.

2 FORMULATION OF THE PROBLEM

2.1 The equations

We consider a reference frame in which an ambient gas of uniform density ρ_a and supersonic speed u_{in} passes through a stationary shock front at $x = x_s$ from upstream i.e. to the right in Fig. 1, as defined by ?. The gas cools and collapses before reaching a cold shell at $x = 0$, also of uniform density. The approach speed u_{in} is defined as a positive quantity. Therefore, the pre-shock gas velocity is $v = -u_{in}$. In this frame, the material in the shell is moving downstream, thus making space for the accumulation of new material.

Following TD93, we take a perpendicular magnetic field, B_a , which is frozen into the gas and so increases in proportion to the density. The shock Alfvén (Mach) number is defined as $M_a = u_{in}/v_a$ where v_a is the Alfvén speed in the ambient medium. The magnetic pressure in the flow is then

$$P_B = b\rho^2, \quad (2)$$

where the constant b is

$$b = \frac{B_a^2}{8\pi\rho_a^2} = v_a^2/2, \quad (3)$$

and we define the useful parameter θ (the parameter β of TD93) as

$$\theta = \frac{B_a^2}{8\pi\rho_a u_{in}^2} = 1/(2 M_a^2), \quad (4)$$

which is the ratio of the magnetic pressure to the upstream ram pressure.




Figure 2. A graphical summary of results for the stability. The growth/damping rates, δ_r plotted against α for the fundamental, first and second mode (from left to right), the indicated different cases of γ (from top to bottom), different β (thick: $\beta = 2$; thin; $\beta = 1$) and three magnetic field strengths, indicated by the line styles shown in the top left-hand panel by the shock Alfvén numbers, M_a .

The one dimensional hydrodynamical equations are (e.g. TD93)

$$\frac{\partial \rho}{\partial t} + \frac{\partial(\rho v)}{\partial x} = 0, \quad (5)$$

$$\rho \left(\frac{\partial v}{\partial t} + v \frac{\partial v}{\partial x} \right) + \frac{\partial P}{\partial x} = 0 \quad (6)$$

and

$$\frac{\partial e}{\partial t} + \frac{\partial(ev)}{\partial x} = -\frac{\partial(Pv)}{\partial x} - \Lambda, \quad (7)$$

where Λ is the radiated energy loss per unit volume and P is the total (magnetic+thermal) pressure related to the internal energy by

$$e = \frac{P}{\gamma - 1} + \left(1 - \frac{1}{\gamma - 1} \right) \frac{B^2}{8\pi} + \frac{1}{2} \rho v^2. \quad (8)$$

Equations (5), (6) & (7) refer to the conservation of mass,

momentum and energy, respectively. Eliminating e ,

$$\frac{\partial P}{\partial t} + v \frac{\partial P}{\partial x} + [\gamma P + b\rho^2(2 - \gamma)] \frac{\partial v}{\partial x} = -(\gamma - 1)A\rho^{\beta - \alpha}(P - b\rho^2)^\alpha, \quad (9)$$

where A is a constant and we have used the ideal gas law to eliminate temperature.

2.2 The steady state

The steady state solution is denoted by the subscript 0. Equations (5) and (6) are integrated to yield

$$\rho_0 v_0 = -\rho_a u_{in} \quad (10)$$

$$P_0 = \rho_a u_{in}(v_0 + u_{in}) + b\rho_a^2 \quad (11)$$

Equations (10) & (11) are substituted in equation (9) which results in

$$\frac{dv_0}{dx} = \frac{C(-\rho_a u_{in})^{\beta - 1} \left[-v_0^2 - u_{in}v_0 + \rho_a b \left(\frac{u_{in}}{v_0} - \frac{v_0}{u_{in}} \right) \right]^\alpha}{v_0^\beta \left[v_0 + \gamma(v_0 + u_{in}) + \rho_a b \left(\frac{\gamma}{u_{in}} + \frac{u_{in}(2 - \gamma)}{v_0^2} \right) \right]}, \quad (12)$$




Figure 3. A graphical summary of results for the frequency. The frequency, δ_i , plotted here against α has been normalised by the factor $1/(2n+1)$ where n is the mode number (fundamental being $n=0$), which demonstrates that the modes resemble that of a half-open organ pipe. Displayed are results for the fundamental, first and second mode (from left to right), the indicated different cases of γ (from top to bottom), different β (thick: $\beta = 2$; thin: $\beta = 1$) and three magnetic field strengths, indicated by the line styles shown in the top left-hand panel by the shock Alfvén numbers, M_a .

where

$$C = (\gamma - 1) * A.$$

This equation can be integrated to determine the velocity through the shocked layer. We now introduce the following variables:

$$\xi = \frac{x}{x_s} \quad (13)$$

$$w = \frac{v_0}{u_{in}}. \quad (14)$$

Equations (12), (13) and (14) lead to

$$\frac{d\xi}{dw} = \frac{-(-w)^\beta \left[w + \gamma(1 + w + \theta) + \frac{(2-\gamma)\theta}{w^2} \right]}{u_{in}^{2\alpha-3} C \rho_a^{\beta-1} x_s \left(-w - w^2 - \theta w + \frac{\theta}{w} \right)^\alpha}. \quad (15)$$

The jump condition across the steady shock front can be

written as (?)

$$\rho_0(x_s) = \rho_a q, \quad (16)$$

$$v_0(x_s) = -\frac{u_{in}}{q}, \quad (17)$$

$$P_0(x_s) = \rho_a u_{in}^2 \left(1 - \frac{1}{q} + \theta \right) \quad (18)$$

where the shock front compression is

$$q = \frac{(U + V)}{4\theta(2 - \gamma)}, \quad (19)$$

$$U = -(\gamma - 1 + 2\gamma\theta),$$

$$V = \left[\sqrt{4\gamma\theta(\gamma\theta - (\gamma - 1)) + (\gamma - 1)^2 + 16\theta} \right].$$

The boundary condition at $x = 0$ is determined from the fact that the temperature is zero: the total pressure is equal to the magnetic pressure and the density remains finite:

$$P_0(x=0) = P_B(x=0). \quad (20)$$

If we denote the velocity of the shell as $v_0(x=0)$, and also




Figure 4. The loci of neutral stability for the indicated values of γ , modes and $\beta = 2$ represented by thick lines and $\beta = 1$ by thin lines. $\log(M_a)$ is plotted against α . The region of instability is above the lines in all cases.

define $w_f = \frac{v_0(x=0)}{u_{in}}$, we can then write from (2) and (11) the expression

$$\rho_a u_{in}^2 (1 + w_f + \theta) = \rho_a u_{in}^2 \left(\frac{\theta}{w_f^2} \right). \quad (21)$$

The physical solution to the above cubic equation is

$$w_f = \frac{-\theta - \sqrt{\theta^2 + 4\theta}}{2}. \quad (22)$$

Note that $-1/w_f$ is the total compression, as indicated in Fig. 1.

2.3 The set of linear equations

The shock wave is now perturbed by

$$\frac{dx_s}{dt} = v_{s1} e^{\sigma t}, \quad (23)$$

where $\sigma = \sigma_R + i\sigma_I$ is the frequency and v_{s1} is a real quantity. The position of the shock may be represented as the real part




Figure 5. The growth rate for various overtones in the presence of a strong transverse magnetic field for the cases $\alpha = 0.3$ (upper panel) and $\alpha = -0.5$ (lower panel) on taking $\beta = 2$, $\gamma = 5/3$ and $M_a = 1.5$.

of

$$x_s = x_{s0} + x_{s1} e^{\sigma t} \quad (24)$$

where $x_{s1} = v_{s1}/\sigma$. Considering only the terms up to first order:

$$\xi = \frac{x}{x_s} = \frac{x}{x_{s0}} \left(1 - \frac{x_{s1}}{x_{s0}} e^{\sigma t} \right) \quad (25)$$

$$\frac{\partial \xi}{\partial x} = \frac{1}{x_{s0}} \left(1 - \frac{x_{s1}}{x_{s0}} e^{\sigma t} \right) \quad (26)$$

$$\frac{\partial \xi}{\partial t} = -\frac{x x_{s1} \sigma e^{\sigma t}}{x_{s0}^2} \quad (27)$$

$$\rho = \rho_0(\xi) + \rho_1(\xi) e^{\sigma t} \quad (28)$$

$$P = P_0(\xi) + P_1(\xi) e^{\sigma t} \quad (29)$$

$$v = v_0(\xi) + v_1(\xi) e^{\sigma t}. \quad (30)$$

All the quantities with subscript 1 represent the small perturbed factors.

When the shock is moving, we consider the frame of reference in which the shock is stationary, i.e., the upstream velocity is now $-u_{in} - v_{s1} e^{\sigma t}$. The boundary conditions at the moving shock wave are obtained by taking the derivatives of equations (16), (17) and (18) with respect to u_{in} to

obtain

$$\rho_1(x = x_s) = 2v_{s1}\theta q^2 \frac{\rho_a}{u_{in}} \frac{d(1/q)}{d\theta}, \quad (31)$$

$$P_1(x = x_s) = 2v_{s1}\rho_a u_{in} \left(1 - \frac{1}{q} + \theta \frac{d(1/q)}{d\theta}\right), \quad (32)$$

$$v_1(x = x_s) = v_{s1} \left(2\theta \frac{d(1/q)}{d\theta} - \frac{1}{q}\right) + v_{s1}. \quad (33)$$

where the last term in (33) arises because of the transformation of coordinates to the shock frame.

We will also need the derivative of q which is, from equation (19),

$$\begin{aligned} \frac{d(1/q)}{d\theta} &= \left[\gamma + \frac{4 + \gamma(2\gamma\theta - (\gamma - 1))}{\sqrt{V}} \right] \\ &\times \left[\frac{2 - \gamma}{2 - \gamma(\gamma - 1)} \right]. \end{aligned} \quad (34)$$

We then transform to the following variables:

$$\zeta = \frac{x_{s0}\sigma\rho_1}{v_{s1}\rho_a}, \quad (35)$$

$$\pi = \frac{P_1}{v_{s1}\rho_a u_{in}}, \quad (36)$$

$$\eta = \frac{v_1}{v_{s1}}, \quad (37)$$

$$\delta = \frac{x_{s0}\sigma}{u_{in}}. \quad (38)$$

At the shock front, from (31) - (33), the new variables take the following values:

$$\xi(w = w_s) = 1, \quad (39)$$

$$\zeta(w = w_s) = 2\delta\theta q^2 \frac{d(1/q)}{d\theta}, \quad (40)$$

$$\pi(w = w_s) = 2 \left(1 - \frac{1}{q} + \theta \frac{d(1/q)}{d\theta}\right), \quad (41)$$

$$\eta(w = w_s) = \left(2\theta \frac{d(1/q)}{d\theta} - \frac{1}{q} + 1\right) \quad (42)$$

Substituting (28), (29) and (30) into (5), (6) and (9), the fluid equations can be rewritten as

$$-\frac{\xi}{w^2} + \zeta \frac{d\xi}{dw} + \frac{w}{\delta} \frac{d\zeta}{dw} + \frac{\eta}{w^2} - \frac{1}{w} \frac{d\eta}{dw} + \frac{\zeta}{\delta} = 0, \quad (43)$$

$$-\xi + \delta \frac{d\zeta}{dw} \eta + w \frac{d\eta}{dw} + \eta - \frac{w^2}{\delta} \zeta - w \frac{d\pi}{dw} = 0, \quad (44)$$

$$\text{where} \quad D + E = F \quad (45)$$

$$\begin{aligned} D &= \left[-\xi + \delta \frac{d\xi}{dw} \pi + w \frac{d\pi}{dw} + \eta - \frac{w}{\delta} + \pi\gamma \right] \\ E &= \left(-\frac{\xi}{w^2} + \zeta \frac{d\xi}{dw} + \frac{w}{\delta} \frac{d\zeta}{dw} + \frac{\eta}{w^2} - \frac{1}{\delta w} + \frac{\zeta}{\delta} \right) \\ &\times \left(\gamma(w + 1 + \theta) + \frac{(2 - \gamma)\theta}{w^2} \right) w - \frac{2(2 - \gamma)\zeta\theta}{w\delta} \\ F &= \left[w + \gamma(1 + w + \theta) + \frac{(2 - \gamma)\theta}{w^2} \right] \\ &\times \left[\alpha\pi - (\beta - \alpha)(1 + w + \theta) \frac{\zeta w}{\delta} + (\beta + \alpha) \frac{\zeta\theta}{\delta w} \right] \\ &\times \left[w + 1 + \theta - \frac{\theta}{w^2} \right]^{-1}. \end{aligned}$$

The quantities ζ , π and η are complex eigenfunctions. We

employ the subscript r to denote the real component and i stands for the imaginary part for each of the above quantities. The quantity δ is a complex number with the sign of the real part, δ_r , indicating the instability (+ve value) or stability (-ve value) of a mode. The quantity δ_i is interpreted as the eigenfrequency (in units of (u_{in}/x_{s0})). The equations for the real and complex quantities are provided in Appendix A.

2.4 The integration process

While integrating the above equations we find that there are numerical problems when $\alpha \leq 0.2$ as the temperature approaches zero. To solve this, we follow a prescription suggested by TD93 which is to introduce a break in the power law cooling at a temperature T_c , at a velocity w_c , leaving a small region near the shell where the cooling function will have $\alpha = 0.5$. Therefore, the cooling function is split into two regimes such that

$$\Lambda = A\rho^{\beta-\alpha} (P - b\rho^2)^\alpha \quad \text{at } w < w_c \quad (46)$$

$$\Lambda = A_c\rho^{\beta-\alpha_c} (P - b\rho^2)^{\alpha_c} \quad \text{at } w > w_c \quad (47)$$

Here, $A_c = A(P_c - b\rho_c)^{\alpha-\alpha_c}$ and we choose $A_c = 0.001$ to be consistent with the results of TD93. To obtain the shock length, we integrate equation (15) from $\xi = 1$ to $\xi = 0$ and take into account the two component cooling function.

The boundary condition at the shell for a moving shock under the influence of a transverse magnetic field is derived in Appendix B. It is more complicated than in Paper I where the gas settles on the wall. In the present scenario, the velocity of the gas does not vanish since the magnetic pressure limits the compression. The boundary condition is

$$|\pi \sqrt{-\frac{w^3}{2\beta}} + \eta| = 0, \quad (48)$$

similar to TD93 but correcting a typographical error in their equation (A27).

For the various combinations of the four free parameters, α , β , γ and θ , the quantities δ_r and δ_i are determined by imposing the boundary condition at the shell boundary. We solved the differential equations employing a fourth order Runge-Kutta technique for trial values of δ_r and δ_i selected from a grid of points uniformly covering the complex plane. The combinations that come closest to satisfying the boundary condition for each mode determine a new set of grid points with a higher resolution.

3 GENERAL RESULTS

We have calculated the growth rates (δ_r) and eigenfrequencies (δ_i) for the various cases of α , β and γ with weak ($M_a = 30$), intermediate ($M_a = 10$) and strong ($M_a = 3$) magnetic fields for the fundamental mode as well as the first three overtones. These results are presented in Tables C1 – C8. Equivalent tables for the case $\gamma = 10/7$ have been constructed but are not presented here. We plot the growth rates against α in Fig. 2 for various modes. The major results are as follows.

- We have accurately reproduced the results of TD93 for the specific case $\beta = 2$ and $\gamma = 5/3$ (Tables C2 & Table C6) as a specific case in our parametric space.

- The fact that the instability regime becomes increasingly restricted with an increase in the transverse magnetic field holds for all values of α , β and γ .

- The fundamental remains the most restricted mode to be unstable. The field has a similar stabilising effect on all the modes.

- The lower value for the density index β provides more stability. However, for high magnetic fields (solid lines in Fig. 2), the stability criterion becomes independent of β . This is interpreted as the cushion effect of the field which limits the compression.

- With decreasing γ , the instability regime is more restricted for all magnetic field strengths. Whether the molecular gas is diatomic ($\gamma = 7/5$) or contains helium ($\gamma = 10/7$) does not significantly alter the instability regime.

- The frequency of the fundamental decreases strongly as α decreases for all magnetic field strengths. The frequency of the overtones generally tend to a constant value as α decreases. Note that in Fig. 3 the frequencies of the modes have been normalised by a factor of $2n + 1$ where n is the mode number for display purposes and so demonstrates that, for all magnetic field strengths, the resonant frequencies are analogous to the acoustical modes in a half-open organ pipe.

The loci of neutral stability in the $\log(M_a) - \alpha$ space, which separate stable and unstable regimes (for fixed β and γ), are displayed in Fig. 4. Whereas an Alfvén number of 8 will suppress the growth of all modes of oscillation for $\alpha > 0$ in the atomic shocks ($\gamma = 5/3, \beta = 2$), we require just $M_a < 20$ for the equivalent molecular case. Hence molecular shocks tend to be considerably more stable. On the other hand, the Alfvén number of 8 would be sufficient to stabilise even the first overtone for molecular shocks with $\alpha = -0.5$.

Given the trends in Fig. 4, it is not clear that higher order modes possess wider instability ranges. In fact, from inspection of the tables (in the appendix), the growth rate tends to be lower for higher overtones for $\alpha > 0.0$ especially as the magnetic field strength increases, whereas for $\alpha < 0$, however, there is an increasing trend in the growth rate. Here, we have investigated these trends for low M_a shocks to determine if the magnetic field will stabilise shocks entirely. As shown in Fig. 5, the growth rate continues to decrease as the mode number is increased for $\alpha = 0.3$ and $M_a = 1.5$, taking an atomic shock for illustration. In contrast, the growth rate displays the opposite trend for the case with negative α . In other words, the magnetic field may not provide complete stability for cooling with negative α . However, the significance of high order modes are probably highly restricted and a multi-dimensional analysis is necessary to determine any observational consequences.

The present analysis applies to both atomic and molecular shocks and assumes that the magnetic field is parallel to the shock front. However, in a sufficiently strong shock the molecules will dissociate immediately following the shock jump. In fact, dissociative cooling may dominate the cooling function. Given a molecular medium with Alfvén speed of 1 km s^{-1} (?) and a speed limit for molecular shocks of 40 km s^{-1} at low densities (under $\sim 10^4 \text{ cm}^{-3}$) and $\sim 24 \text{ km s}^{-1}$ at high cloud densities (?), the maximum Alfvén number for molecular shocks with $\gamma = 7/5$ is then in the range 24 – 40.

A maximum value for the magnetic field should also exist, above which ambipolar diffusion dominates the shock

physics. The limit is more difficult to calculate since it depends on the ionisation fraction, density and cooling function. In fact, the limitation is much stronger on the ion fraction rather than the magnetic field. In molecular clouds which are well shielded from the extreme ultraviolet, the ion fraction, χ , is extremely low and ambipolar diffusion dominates. On the other hand, the ion fraction may reach high values near massive stars, stellar outflows or within bow shocks with high-speed apices, supplying ionising radiation. As estimated by ?, ion fractions exceeding $\sim 10^{-5} B_{-3}/n_6^{3/2}$ (H_2O cooling) or $\sim 2 \cdot 10^{-6} B_{-3}/n_6^{1/2}$ (H_2 cooling) are required to ensure a J-type shock where $B_{-3} = B/(10^{-3} \text{ G})$ and $n_6 = n/(10^6 \text{ cm}^{-3})$. Thus, the Alfvén speed must not exceed $\sim 5 \cdot 10^5 n_6 \chi$ (H_2O cooling) km s^{-1} (H_2O cooling) or $\sim 10^5 \chi \text{ km s}^{-1}$ (H_2 cooling).

4 CONCLUSIONS: MOLECULAR SHOCK

We have found the stable and unstable regions for radiative shocks in the α , β and γ parameter space for various magnetic field strengths and in the field- α parameter space for specific β and γ values.

For the fundamental mode, molecular shocks are considerably more stable than atomic shocks. For the overtones, molecular shocks are only moderately more stable. The magnetic field, however, has a strong stabilising influence on all these modes, decreasing all the growth rates by similar factors (see Fig. 2).

The field strength enters the stability condition through the shock Alfvén number. All cooling functions in which the cooling increases with temperature are stable for $M_a < 20$. However, for $M_a > 22$, the first overtone can be unstable, potentially leading to oscillations with non-linear amplitudes. Hence, we need to estimate the Alfvén number for typical shocks in various environments.

Unfortunately, in molecular clouds, the Alfvén speed is a very uncertain parameter. It is even difficult to get an objective view on the magnetic field strength alone (?) although derived Alfvén speeds generally lie within the range $0.5 - 5 \text{ km s}^{-1}$ (?). Jump shocks followed by cooling zones can arise without molecular dissociation for shock speeds as high as 40 km s^{-1} at low densities but $\sim 24 \text{ km s}^{-1}$ at high cloud densities (?). As discussed in § 3, a minimum ion fraction is also required to maintain a frozen-in magnetic field. Hence, a quite fast shock in a diffuse molecular cloud is the most favourable state for the appearance of oscillations.

The temperature dependence of molecular cooling functions generally correspond to values of α well above zero and so indicates stable shocks. However, besides temperature and density, shock cooling may depend on the chemistry. The rate of formation of trace molecules within the cooling layer can increase the cooling rate as the temperature falls. In particular, the propagation of warming but non-ionising shocks into cool atomic gas may show trace molecule formation and increasing cooling within a gas with $\gamma = 5/3$. However, because of the low shock speeds, these shocks would be stabilised if the magnetic field were transverse. Hence, it can be that oblique non-ionising shock waves into atomic gas represent the rather restrictive conditions for shock instability within the cool components of the interstellar medium.

Although it is well known how an oblique field alters

the nature of a steady shock, it is not clear how it alters the stability conditions. It should also be noted that the magnetic field may help stabilise shocks even when the field is parallel to the shock provided the Alfvén number is less than the Mach number, a condition expected to be satisfied in molecular clouds (?). Such an analysis, as well as multi-dimensional numerical simulations of radiative shocks, are still to be performed.

ACKNOWLEDGMENTS

Research at Armagh Observatory is funded by the Department of Culture, Arts and Leisure, Northern Ireland. BR is extremely grateful to Sathya Sai Baba for inspiration and also thanks Professor S. Kandaswami for encouragement and advice.

APPENDIX A: THE DIFFERENTIAL EQUATIONS

From Eqs. (36), (37) and (38), we get six coupled first order equations which are

$$\begin{aligned} \frac{d\eta_r}{dw} &= \frac{\alpha\pi_r\delta^2 + (\delta_r\zeta_r + \delta_i\zeta_i) \left(\frac{(\beta+\alpha)\theta}{w} - w(\beta-\alpha)(1+w+\theta) \right)}{(1+w+\theta - \frac{\theta}{w^2})\delta^2} \\ &+ \frac{2\xi\delta^2 - 2\eta_r\delta^2 - \gamma\pi_r\delta^2 + (w^2 + \frac{2(2-\gamma)\theta}{w})(\zeta_r\delta_r + \zeta_i\delta_i)}{(w + \gamma(w+1+\theta) + \frac{(2-\gamma)\theta}{w^2})\delta^2} \\ &- \frac{d\xi}{dw} \left[\frac{\delta_r(\eta_r + \pi_r) - \delta_i(\eta_i + \pi_i)}{(w + \gamma(w+1+\theta) + \frac{(2-\gamma)\theta}{w^2})} \right] + \frac{\delta_r}{\delta^2} \quad (A1) \\ \frac{d\eta_i}{dw} &= \frac{\alpha\pi_i\delta^2 + (\delta_r\zeta_i - \delta_i\zeta_r) \left(\frac{(\beta+\alpha)\theta}{w} - w(\beta-\alpha)(1+w+\theta) \right)}{(1+w+\theta - \frac{\theta}{w^2})\delta^2} \\ &+ \frac{-2\eta_i\delta^2 - \gamma\pi_i\delta^2 + (w^2 + \frac{2(2-\gamma)\theta}{w})(\zeta_i\delta_r - \zeta_r\delta_i)}{(w + \gamma(w+1+\theta) + \frac{(2-\gamma)\theta}{w^2})\delta^2} \\ &- \frac{d\xi}{dw} \left[\frac{\delta_i(\eta_r + \pi_r) + \delta_r(\eta_i + \pi_i)}{(w + \gamma(w+1+\theta) + \frac{(2-\gamma)\theta}{w^2})} \right] - \frac{\delta_i}{\delta^2} \\ \frac{d\pi_r}{dw} &= \frac{-\xi}{w} + \frac{(\delta_r\eta_r - \delta_i\eta_i)}{w} \frac{d\xi}{dw} + \frac{\eta_r}{w} + \frac{d\eta_r}{dw} \\ &- \frac{w(\zeta_r\delta_r + \zeta_i\delta_i)}{\delta^2} \quad (A3) \end{aligned}$$

$$\begin{aligned} \frac{d\pi_i}{dw} &= \frac{(\delta_i\eta_r + \delta_r\eta_i)}{w} \frac{d\xi}{dw} + \frac{\eta_i}{w} + \frac{d\eta_i}{dw} \\ &- \frac{w(\zeta_i\delta_r - \zeta_r\delta_i)}{\delta^2} \quad (A4) \end{aligned}$$

$$\begin{aligned} \frac{d\zeta_r}{dw} &= \frac{\xi\delta_r}{w^3} - \frac{d\xi}{dw} \frac{\zeta_r\delta_r}{w} + \frac{\delta_r}{w^2} \frac{d\eta_r}{dw} - \frac{\eta_r\delta_r}{w^3} \\ &- \frac{\delta_r(\zeta_r\delta_r + \zeta_i\delta_i)}{w\delta^2} + \frac{d\xi}{dw} \frac{\zeta_i\delta_i}{w} - \frac{\delta_i}{w^2} \frac{d\eta_i}{dw} \\ &+ \frac{\eta_i\delta_i}{w^3} + \frac{\delta_i(\zeta_i\delta_r - \zeta_r\delta_i)}{w\delta^2} \quad (A5) \end{aligned}$$

$$\begin{aligned} \frac{d\zeta_i}{dw} &= \frac{\xi\delta_i}{w^3} - \frac{d\xi}{dw} \frac{\zeta_i\delta_r}{w} + \frac{\delta_r}{w^2} \frac{d\eta_i}{dw} - \frac{\eta_i\delta_r}{w^3} \\ &- \frac{\delta_r(\zeta_i\delta_r - \zeta_r\delta_i)}{w\delta^2} - \frac{d\xi}{dw} \frac{\zeta_r\delta_i}{w} + \frac{\delta_i}{w^2} \frac{d\eta_r}{dw} \\ &- \frac{\eta_r\delta_i}{w^3} - \frac{\delta_i(\zeta_r\delta_r + \zeta_j\delta_i)}{w\delta^2}. \quad (A6) \end{aligned}$$

APPENDIX B: BOUNDARY CONDITION AT THE SHELL

Following TD93, we exploit the condition that the temperature tends to zero at the shell. This implies that the only pressure at the shell is due to the magnetic field. To extract the boundary condition, we represent the physical quantities as

$$\rho = \rho_0(x) + \rho_1(x)e^{i(ft+kx)} \quad (B1)$$

$$P = P_0(x) + P_1(x)e^{i(ft+kx)} \quad (B2)$$

$$v = v_0(x) + v_1(x)e^{i(ft+kx)} \quad (B3)$$

where k is the wavenumber, which represents the direction of the outgoing wave towards the shell and f is the frequency. The above equations are substituted in (5) and (6) to yield

$$\frac{f}{k} = - \left(\frac{\rho_0 v_1 v_0 + P_1}{\rho_0 v_1} \right) \quad (B4)$$

$$\frac{f}{k} = - \left(\frac{\rho_0 v_1 + \rho_1 v_0}{\rho_1} \right) \quad (B5)$$

Equating the above expressions leads to

$$P_1 = \frac{\rho_0^2 v_1^2}{\rho_1} \quad (B6)$$

The magnetic pressure is given by

$$P = \frac{B^2}{8\pi} \quad (B7)$$

We now express the magnetic field in terms of density for example, (?) as

$$\frac{B}{B_a} = \frac{\rho}{\rho_a} \quad (B8)$$

(A2) where B and ρ are the post-shock magnetic field and density. Substituting (B2) in (B7) and defining the Alfvén velocity as $v_A = \sqrt{\frac{B^2}{4\pi\rho}}$, we get

$$P_1 = v_A^2 \rho_1 \quad (B9)$$

Equations (B6) and (B9) yield the boundary condition at the shell as,

$$P_1 = -v_A v_1 \rho_0 \quad (B10)$$

The minus sign indicates the direction of the gas flow. Note that the perturbed quantities are function of x , which itself is a function of ξ from (25). The expression (B10) can be

written in terms of the non-dimensional quantities as

$$\pi = -\eta \sqrt{\frac{2\theta}{-w_f^3}} \quad (\text{B11})$$

The above quantities are evaluated at the shell boundary or equivalently at $w = w_f$.

APPENDIX C: TABLES

Table C1. Growth rates (δ_r) for several modes and magnetic field strengths, as measured by the Alfvén number M_a , for $\gamma = 5/3$, $\beta = 1$.

fundamental				first overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	-0.02942	0.04903	0.08965	0.06387	0.11420	0.14986
-0.5	-0.08446	0.00567	0.03646	-0.03742	0.03890	0.07482
0.0	-0.18538	-0.05842	-0.02296	-0.20521	-0.06762	-0.01974
0.3	-0.29720	-0.11769	-0.07122	-0.38342	-0.16282	-0.09245
0.5	-0.42465	-0.17862	-0.11462	-0.57743	-0.25705	-0.15414

second overtone				third overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	0.11050	0.13937	0.16695	0.13817	0.16075	0.19142
-0.5	-0.00862	0.04990	0.07818	0.00809	0.06541	0.09936
0.0	-0.20728	-0.07856	-0.03713	-0.21087	-0.07366	-0.02433
0.3	-0.42137	-0.19620	-0.12957	-0.45058	-0.20419	-0.12893
0.5	-0.65377	-0.31342	-0.21165	-0.70913	-0.33717	-0.22760

Table C2. Growth rates (δ_r) for several modes and magnetic field strengths for $\gamma = 5/3$, $\beta = 2$. This Table can be compared to Table 1 of TD93.

fundamental				first overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	-0.03549	0.04610	0.09351	0.06696	0.13981	0.19781
-0.5	-0.08877	0.01342	0.04383	-0.02921	0.08413	0.13291
0.0	-0.17983	-0.02853	0.00702	-0.18560	0.00496	0.06983
0.3	-0.27889	-0.06263	-0.01806	-0.35272	-0.06229	0.02479
0.5	-0.39054	-0.09283	-0.03697	-0.53889	-0.12623	-0.00998

second overtone				third overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	0.11374	0.17466	0.22738	0.14253	0.19826	0.25109
-0.5	0.00067	0.10719	0.16104	0.01897	0.12425	0.18399
0.0	-0.18654	0.00890	0.08662	-0.18846	0.01377	0.10306
0.3	-0.38992	-0.08046	0.03131	-0.41714	-0.09122	0.04081
0.5	-0.61605	-0.17289	-0.01352	-0.67020	-0.20545	-0.01179

Table C3. Growth rates (δ_r) for several modes and magnetic field strengths for $\gamma = 7/5$, $\beta = 1$.

fundamental				first overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	-0.12574	-0.03009	0.00029	-0.04055	0.02876	0.06090
-0.5	-0.18312	-0.06286	-0.02743	-0.13786	-0.02838	0.01337
0.0	-0.28865	-0.11479	-0.06609	-0.30446	-0.11369	-0.05206
0.3	-0.40691	-0.16801	-0.10022	-0.48460	-0.19620	-0.10778
0.5	-0.53663	-0.22589	-0.13324	-0.67452	-0.28096	-0.16065

second overtone				third overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	0.00362	0.04071	0.05939	0.02880	0.06460	0.08899
-0.5	-0.11048	-0.02764	0.00114	-0.09556	-0.09275	0.02439
0.0	-0.30800	-0.13002	-0.07824	-0.31358	-0.12335	-0.06976
0.3	-0.52493	-0.22824	-0.14361	-0.55789	-0.23731	-0.15350
0.5	-0.74988	-0.32688	-0.20005	-0.80390	-0.35412	-0.22737

Table C4. Growth rates (δ_r) for several modes and magnetic field strengths for $\gamma = 7/5$, $\beta = 2$.

fundamental				first overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	-0.12850	-0.02761	0.01627	-0.03781	0.04672	0.08856
-0.5	-0.18572	-0.04914	-0.01205	-0.13274	0.00162	0.05346
0.0	-0.28557	-0.08540	-0.03462	-0.29240	-0.06666	0.01054
0.3	-0.39682	-0.11950	-0.05177	-0.46628	-0.13231	-0.02255
0.5	-0.51900	-0.15418	-0.06567	-0.65304	-0.20231	-0.05032
second overtone				third overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	0.00628	0.06857	0.10574	0.03269	0.09028	0.13057
-0.5	-0.10479	0.01285	0.06522	-0.08846	0.02882	0.08690
0.0	-0.29458	-0.07317	0.01163	-0.29851	-0.06802	0.02653
0.3	-0.50449	-0.15932	-0.03224	-0.53521	-0.16798	-0.02552
0.5	-0.72683	-0.25388	-0.07193	-0.77992	-0.27967	-0.07602

Table C5. Eigenfrequencies (δ_i) for several modes and magnetic field strengths for $\gamma = 5/3$ and $\beta = 1$.

fundamental				first overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	0.44847	0.33935	0.30432	1.45520	1.03328	0.92059
-0.5	0.50430	0.37630	0.34126	1.50546	1.03500	0.91030
0.0	0.58627	0.41387	0.36464	1.59283	1.03047	0.87269
0.3	0.67073	0.44723	0.37841	1.68773	1.02950	0.83627
0.5	0.75300	0.48020	0.38821	1.76664	1.02936	0.80303
second overtone				third overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	2.50463	1.77852	1.59824	3.55657	2.51123	2.25140
-0.5	2.54768	1.74493	1.53713	3.59392	2.44672	2.14906
0.0	2.63290	1.69280	1.43223	3.67699	2.35191	1.98756
0.3	2.72930	1.65092	1.33791	3.77286	2.27393	1.84544
0.5	2.79538	1.60990	1.24986	3.82144	2.19758	1.71171

Table C6. Eigenfrequencies (δ_i) for several modes and magnetic field strengths for $\gamma = 5/3$ and $\beta = 2$.

fundamental				first overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	0.43031	0.27658	0.25635	1.45075	1.03787	0.94698
-0.5	0.47773	0.32531	0.28242	1.49626	1.05480	0.95981
0.0	0.55034	0.35176	0.30317	1.58012	1.06036	0.94929
0.3	0.62587	0.37345	0.31234	1.67378	1.07176	0.93804
0.5	0.70256	0.39530	0.31871	1.75753	1.09095	0.92967
second overtone				third overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	2.50320	1.81950	1.68275	3.55754	2.59039	2.40284
-0.5	2.54284	1.80897	1.66530	3.59230	2.55669	2.36004
0.0	2.62573	1.78459	1.62070	3.67348	2.50237	2.28419
0.3	2.72132	1.77329	1.58450	3.76880	2.46542	2.22368
0.5	2.79310	1.77345	1.55688	3.82473	2.43991	2.17669

Table C7. Eigenfrequencies (δ_i) for several modes and varying magnetic field strengths for $\gamma = 7/5$ and $\beta = 1$.

fundamental				first overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	0.43865	0.27943	0.23392	1.41341	0.84427	0.69527
-0.5	0.48338	0.30069	0.25072	1.46172	0.84617	0.68237
0.0	0.55590	0.32986	0.26344	1.55387	0.85161	0.65657
0.3	0.62987	0.35788	0.27225	1.64946	0.86141	0.63397
0.5	0.69568	0.38442	0.27884	1.71604	0.86861	0.61193
second overtone				third overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	2.43144	1.44978	1.19709	3.45026	2.04889	1.69333
-0.5	2.47796	1.42644	1.14496	3.49517	2.00664	1.61494
0.0	2.57765	1.39673	1.06321	3.60095	1.95155	1.49445
0.3	2.68125	1.37676	0.99033	3.70963	1.90977	1.38357
0.5	2.73264	1.35376	0.92053	3.74108	1.85977	1.27176

Table C8. Eigenfrequencies (δ_i) for the various modes of parameters $\gamma = 7/5$ and $\beta = 2$ for various magnetic fields.

fundamental				first overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	0.42613	0.24811	0.19424	1.41061	0.84868	0.71088
-0.5	0.46608	0.26937	0.21581	1.45638	0.85532	0.71608
0.0	0.53258	0.29198	0.22668	1.54505	0.86521	0.70578
0.3	0.60136	0.31426	0.23328	1.63895	0.88205	0.69826
0.5	0.66468	0.33806	0.23891	1.70814	0.90367	0.69442
second overtone				third overtone		
α	$M_a = 3$	$M_a = 10$	$M_a = 30$	$M_a = 3$	$M_a = 10$	$M_a = 30$
-1	2.42930	1.47335	1.25896	3.44982	2.09150	1.79406
-0.5	2.47396	1.46111	1.23842	3.49352	2.06317	1.75504
0.0	2.57015	1.44656	1.20060	3.59567	2.02658	1.69333
0.3	2.67207	1.44242	1.17200	3.70268	2.00303	1.64574
0.5	2.72750	1.44027	1.15139	3.73955	1.97825	1.60978